

Cognitive behavioral therapy (CBT) for adult posttraumatic stress disorder (PTSD) Adult Mental Health: Trauma

Benefit-cost estimates updated December 2019. Literature review updated September 2016.

Current estimates replace old estimates. Numbers will change over time as a result of model inputs and monetization methods.

The WSIPP benefit-cost analysis examines, on an apples-to-apples basis, the monetary value of programs or policies to determine whether the benefits from the program exceed its costs. WSIPP's research approach to identifying evidence-based programs and policies has three main steps. First, we determine "what works" (and what does not work) to improve outcomes using a statistical technique called meta-analysis. Second, we calculate whether the benefits of a program exceed its costs. Third, we estimate the risk of investing in a program by testing the sensitivity of our results. For more detail on our methods, see our [Technical Documentation](#).

Program Description: Treatments in this review include several components, such as psycho-education about posttraumatic stress disorder (PTSD), relaxation and other techniques for managing physiological and emotional stress, exposure (the gradual desensitization to memories of the traumatic event), and cognitive restructuring of inaccurate or unhelpful thoughts. The studies in this review employed a number of trauma-specific treatment models including Prolonged Exposure Therapy (PE), Narrative Exposure Therapy (NET), and Cognitive Processing Therapy (CPT). In the studies in this review, treatments provided between 1-45 therapeutic hours per client in individual or group settings. Studies were conducted on all continents and subjects had experienced one of a variety of types of trauma including terrorism, sexual or physical assault, domestic violence, war, political detention, and automobile accidents.

Benefit-Cost Summary Statistics Per Participant

Benefits to:

Taxpayers	\$16,561	Benefit to cost ratio	\$88.95
Participants	\$29,093	Benefits minus costs	\$52,092
Others	\$4,898	Chance the program will produce	
Indirect	\$2,132	benefits greater than the costs	100 %
Total benefits	\$52,684		
Net program cost	(\$592)		
Benefits minus cost	\$52,092		

The estimates shown are present value, life cycle benefits and costs. All dollars are expressed in the base year chosen for this analysis (2018). The chance the benefits exceed the costs are derived from a Monte Carlo risk analysis. The details on this, as well as the economic discount rates and other relevant parameters are described in our [Technical Documentation](#).

Meta-Analysis of Program Effects

Outcomes measured	Treatment age	No. of effect sizes	Treatment N	Adjusted effect sizes and standard errors used in the benefit-cost analysis						Unadjusted effect size (random effects model)	
				First time ES is estimated			Second time ES is estimated			ES	p-value
				ES	SE	Age	ES	SE	Age		
Anxiety disorder	39	17	355	-0.620	0.087	40	-0.620	0.087	41	-0.948	0.001
Employment^^	39	1	12	0.348	0.530	40	n/a	n/a	n/a	0.821	0.125
Major depressive disorder	39	49	1389	-0.433	0.046	40	-0.433	0.046	41	-0.717	0.001
Post-traumatic stress	39	70	2361	-0.539	0.047	40	-0.539	0.047	41	-0.950	0.001
Substance use disorder^	39	1	55	-0.164	0.366	40	n/a	n/a	n/a	-0.261	0.477

^ WSIPP's benefit-cost model does not monetize this outcome.

^^ WSIPP does not include this outcome when conducting benefit-cost analysis for this program.

Meta-analysis is a statistical method to combine the results from separate studies on a program, policy, or topic in order to estimate its effect on an outcome. WSIPP systematically evaluates all credible evaluations we can locate on each topic. The outcomes measured are the types of program impacts that were measured in the research literature (for example, crime or educational attainment). Treatment N represents the total number of individuals or units in the treatment group across the included studies.

An effect size (ES) is a standard metric that summarizes the degree to which a program or policy affects a measured outcome. If the effect size is positive, the outcome increases. If the effect size is negative, the outcome decreases.

Adjusted effect sizes are used to calculate the benefits from our benefit cost model. WSIPP may adjust effect sizes based on methodological characteristics of the study. For example, we may adjust effect sizes when a study has a weak research design or when the program developer is involved in the research. The magnitude of these adjustments varies depending on the topic area.

WSIPP may also adjust the second ES measurement. Research shows the magnitude of some effect sizes decrease over time. For those effect sizes, we estimate outcome-based adjustments which we apply between the first time ES is estimated and the second time ES is estimated. We also report the unadjusted effect size to show the effect sizes before any adjustments have been made. More details about these adjustments can be found in our [Technical Documentation](#).

Detailed Monetary Benefit Estimates Per Participant

Affected outcome:	Resulting benefits: ¹	Benefits accrue to:				
		Taxpayers	Participants	Others ²	Indirect ³	Total
Post-traumatic stress	Labor market earnings associated with PTSD	\$11,813	\$27,747	\$0	\$0	\$39,560
Post-traumatic stress	Health care associated with PTSD	\$4,747	\$1,342	\$4,898	\$2,373	\$13,360
Major depressive disorder	Mortality associated with depression	\$2	\$4	\$0	\$55	\$60
Program cost	Adjustment for deadweight cost of program	\$0	\$0	\$0	(\$296)	(\$296)
Totals		\$16,561	\$29,093	\$4,898	\$2,132	\$52,684

¹In addition to the outcomes measured in the meta-analysis table, WSIPP measures benefits and costs estimated from other outcomes associated with those reported in the evaluation literature. For example, empirical research demonstrates that high school graduation leads to reduced crime. These associated measures provide a more complete picture of the detailed costs and benefits of the program.

²"Others" includes benefits to people other than taxpayers and participants. Depending on the program, it could include reductions in crime victimization, the economic benefits from a more educated workforce, and the benefits from employer-paid health insurance.

³"Indirect benefits" includes estimates of the net changes in the value of a statistical life and net changes in the deadweight costs of taxation.


Detailed Annual Cost Estimates Per Participant

	Annual cost	Year dollars	Summary	
Program costs	\$1,444	2014	Present value of net program costs (in 2018 dollars)	(\$592)
Comparison costs	\$814	2008	Cost range (+ or -)	15 %

These therapies can take place over 1-45 weekly sessions; total length of treatment is less than one year. The per-participant cost of treatment by modality (group/individual) was weighted by the treatment Ns reported in the studies. Cost per session is \$40.04/session for group and \$122.25 for individual therapy (2015 dollars). This rate is based on actuarial tables reported in Mercer (2014) Behavioral Health Data Book for the State of Washington For Rates Effective January 1, 2015. The comparison group costs are from the average Medicaid expenditures for PTSD treatment in Washington in 2009.


The figures shown are estimates of the costs to implement programs in Washington. The comparison group costs reflect either no treatment or treatment as usual, depending on how effect sizes were calculated in the meta-analysis. The cost range reported above reflects potential variation or uncertainty in the cost estimate; more detail can be found in our [Technical Documentation](#).

Benefits Minus Costs Over Time (Cumulative Discounted Dollars)



The graph above illustrates the estimated cumulative net benefits per-participant for the first fifty years beyond the initial investment in the program. We present these cash flows in discounted dollars. If the dollars are negative (bars below \$0 line), the cumulative benefits do not outweigh the cost of the program up to that point in time. The program breaks even when the dollars reach \$0. At this point, the total benefits to participants, taxpayers, and others, are equal to the cost of the program. If the dollars are above \$0, the benefits of the program exceed the initial investment.

Benefits by Perspective Over Time (Cumulative Discounted Dollars)



The graph above illustrates the breakdown of the estimated cumulative benefits (not including program costs) per-participant for the first fifty years beyond the initial investment in the program. These cash flows provide a breakdown of the classification of dollars over time into four perspectives: taxpayer, participant, others, and indirect. "Taxpayers" includes expected savings to government and expected increases in tax revenue. "Participants" includes expected increases in earnings and expenditures for items such as health care and college tuition. "Others" includes benefits to people other than taxpayers and participants. Depending on the program, it could include reductions in crime victimization, the economic benefits from a more educated workforce, and the benefits from employer-paid health insurance. "Indirect benefits" includes estimates of the changes in the value of a statistical life and changes in the deadweight costs of taxation. If a section of the bar is below the \$0 line, the program is creating a negative benefit, meaning a loss of value from that perspective.

Taxpayer Benefits by Source of Value Over Time (Cumulative Discounted Dollars)



The graph above focuses on the subset of estimated cumulative benefits that accrue to taxpayers. The cash flows are divided into the source of the value.

Citations Used in the Meta-Analysis

- Alghamdi, M., Hunt, N., & Thomas, S. (2015). The effectiveness of Narrative Exposure Therapy with traumatised firefighters in Saudi Arabia: A randomized controlled study. *Behaviour Research and Therapy, 66*, 64-71.
- Asukai, N., Saito, A., Tsuruta, N., Kishimoto, J., & Nishikawa, T. (2010). Efficacy of exposure therapy for Japanese patients with posttraumatic stress disorder due to mixed traumatic events: A randomized controlled study. *Journal of Traumatic Stress, 23*(6), 744-750.
- Basoglu, M., Salcioglu, E., & Livanou, M. (2007). A randomized controlled study of single-session behavioural treatment of earthquake-related post-traumatic stress disorder using an earthquake simulator. *Psychological Medicine, 37*(2), 203-13.
- Basoglu, M., Salcioglu, E., Livanou, M., Kalender, D., & Acar, G. (2005). Single-session behavioral treatment of earthquake-related posttraumatic stress disorder: a randomized waiting list controlled trial. *Journal of Traumatic Stress, 18*(1), 1-11.
- Beck, J.G., Coffey, S.F., Foy, D.W., Keane, T.M., & Blanchard, E.B. (2009). Group cognitive behavior therapy for chronic posttraumatic stress disorder: An initial randomized pilot study. *Behavior Therapy, 40*(1), 82-92.
- Bichescu, D., Neuner, F., Schaer, M., Elbert, T. (2007) Narrative exposure therapy for political imprisonment-related chronic posttraumatic stress disorder and depression. *Behaviour Research and Therapy, 45*(9), 2212-2220.
- Brom, D., Kleber, R.J., & Defares, P.B. (1989). Brief psychotherapy for posttraumatic stress disorders. *Journal of Consulting and Clinical Psychology, 57*(5), 607-612.
- Bryant, R.A., Ekasawin, S., Chakrabhand, S., Suwanmitri, S., Duangchun, O., & Chantaluckwong, T. (2011). A randomized controlled effectiveness trial of cognitive behavior therapy for post-traumatic stress disorder in terrorist-affected people in Thailand. *World Psychiatry, 10*(3), 205-209.
- Bryant, R.A., Moulds, M.L., Guthrie, R.M., Dang, S.T., & Nixon, R.D.V. (2003). Imaginal exposure alone and imaginal exposure with cognitive restructuring in treatment of posttraumatic stress disorder. *Journal of Consulting and Clinical Psychology, 71*(4), 706-712.
- Castillo, D.T., Chee, C.L., Nason, E., Keller, J., C'de, B.J., Qualls, C., Fallon, S.K., . . . Keane, T.M. (2016). Group-delivered cognitive/exposure therapy for PTSD in women veterans: A randomized controlled trial. *Psychological Trauma: Theory, Research, Practice, and Policy, 8*(3), 404.
- Chard, K.M. (2005). An evaluation of cognitive processing therapy for the treatment of posttraumatic stress disorder related to childhood sexual abuse. *Journal of Consulting and Clinical Psychology, 73*(5), 965-971.
- Cottraux, J., Note, I., Yao, S.N., de Mey-Guillard, C., Bonasse, F., Djamoussian, D., Mollard, E., . . . Chen, Y. (2008). Randomized controlled comparison of cognitive behavior therapy with Rogerian Supportive Therapy in chronic post-traumatic stress disorder: A 2-year follow-up. *Psychotherapy and Psychosomatics, 77*(2), 101-110.
- Difede, J., Malta, L.S., Best, S., Henn-Haase, C., Metzler, T., Bryant, R., & Marmar, C. (2007). A randomized controlled clinical treatment trial for World Trade Center attack-related PTSD in disaster workers. *The Journal of Nervous and Mental Disease, 195*(10), 861-5.
- Dorrepaal, E., Thomaes, K., Smit, J.H., van Balkom, A.J., Veltman, D.J., Hoogendoorn, A.W., & Draijer, N. (2012). Stabilizing group treatment for complex posttraumatic stress disorder related to child abuse based on psychoeducation and cognitive behavioural therapy: a multisite randomized controlled trial. *Psychotherapy and Psychosomatics, 81*(4), 217-25.
- Duffy, M., Gillespie, K., & Clark, D.M. (2007). Post-traumatic stress disorder in the context of terrorism and other civil conflict in Northern Ireland: randomised controlled trial. *BMJ, 334*(7604), 1147.
- Dunne, R.L., Kenardy, J., & Sterling, M. (2012). A randomized controlled trial of cognitive-behavioral therapy for the treatment of PTSD in the context of chronic whiplash. *The Clinical Journal of Pain, 28*(9), 755-765.
- Echeburua, E., Corral, P.D., Zubizarreta, I., & Sarasua, B. (1997). Psychological treatment of chronic posttraumatic stress disorder in victims of sexual aggression. *Behavior Modification, 21*(4), 433-456.
- Ehlers, A., Clark, D.M., Hackmann, A., McManus, F., Fennell, M., Herbert, C., & Mayou, R. (2003). A randomized controlled trial of cognitive therapy, a self-help booklet, and repeated assessments as early interventions for posttraumatic stress disorder. *Archives of General Psychiatry, 60*(10), 1024-1032.
- Ehlers, A., Clark, D.M., Hackmann, A., McManus, F., & Fennell, M. (2005). Cognitive therapy for post-traumatic stress disorder: development and evaluation. *Behaviour Research and Therapy, 43*(4), 413-431.
- Ehlers, A., Hackmann, A., Grey, N., Wild, J., Liness, S., Albert, I., Deale, A., . . . Clark, D.M. (2014). A randomized controlled trial of 7-day intensive and standard weekly cognitive therapy for PTSD and emotion-focused supportive therapy. *The American Journal of Psychiatry, 171*(3), 294-304.
- Fecteau, G., & Nicki, R. (1999). Cognitive behavioural treatment of post traumatic stress disorder after motor vehicle accident. *Behavioural and Cognitive Psychotherapy, 27*(3), 201-214.
- Feske, U. (2008). Treating low-income and minority women with posttraumatic stress disorder: A pilot study comparing prolonged exposure and treatment as usual conducted by community therapists. *Journal of Interpersonal Violence, 23*(8), 1027-1040.
- Foa, E.B., Rothbaum, B.O., Riggs, D.S., & Murdock, T.B. (1991). Treatment of posttraumatic stress disorder in rape victims: A comparison between cognitive-behavioral procedures and counseling. *Journal of Consulting and Clinical Psychology, 59*(5), 715-723.
- Foa, E.B., Dancu, C.V., Hembree, E.A., Jaycox, L.H., Meadows, E.A., Street, G.P. (1999). A comparison of exposure therapy, stress inoculation therapy, and their combination for reducing posttraumatic stress disorder in female assault victims. *Journal of Consulting and Clinical Psychology, 67*(2), 194-200
- Foa, E.B., Hembree, E.A., Cahill, S.P., Rauch, S.A.M., Riggs, D.S., Feeny, N.C., & Yadin, E. (2005). Randomized trial of prolonged exposure for posttraumatic stress disorder with and without cognitive restructuring: Outcome at academic and community clinics. *Journal of Consulting and Clinical Psychology, 73*(5), 953-964.
- Forbes, D., Lloyd, D., Nixon, R.D.V., Elliott, P., Varker, T., Perry, D., Bryant, R.A., . . . Creamer, M. (2012). A multisite randomized controlled effectiveness trial of cognitive processing therapy for military-related posttraumatic stress disorder. *Journal of Anxiety Disorders, 26*(3), 442-452.
- Gersons, B.P., Carlier, I.V., Lamberts, R.D., & van der Kolk, B.A. (2000). Randomized clinical trial of brief eclectic psychotherapy for police officers with posttraumatic stress disorder. *Journal of Traumatic Stress, 13*(2), 333-347.
- Hinton, D.E., Pollack, M.H., Hofmann, S.G., & Otto, M.W. (2009). Mechanisms of efficacy of CBT for Cambodian refugees with PTSD: Improvement in emotion regulation and orthostatic blood pressure response. *CNS Neuroscience and Therapeutics, 15*(3), 255-263.
- Hinton, D.E., Hofmann, S.G., Rivera, E., Otto, M.W., & Pollack, M.H. (2011). Culturally adapted CBT (CA-CBT) for Latino women with treatment-resistant PTSD: A pilot study comparing CA-CBT to applied muscle relaxation. *Behaviour Research and Therapy, 49*(4), 275-280.
- Hinton, D.E., Chhean, D., Pich, V., Safren, S.A., Hofmann, S.G., & Pollack, M.H. (2005). A randomized controlled trial of cognitive-behavior therapy for Cambodian refugees with treatment-resistant PTSD and panic attacks: a cross-over design. *Journal of Traumatic Stress, 18*(6), 617-29.

- Hollifield, M., Sinclair-Lian, N., Warner, T.D., & Hammerschlag, R. (2007). Acupuncture for posttraumatic stress disorder: a randomized controlled pilot trial. *The Journal of Nervous and Mental Disease, 195*(6), 504-13.
- Johnson, D.M., Zlotnick, C., & Perez, S. (2011). Cognitive behavioral treatment of PTSD in residents of battered women's shelters: Results of a randomized clinical trial. *Journal of Consulting and Clinical Psychology, 79*(4), 542-551.
- Kent, M., Davis, M.C., Stark, S.L., & Stewart, L.A. (2011). A resilience-oriented treatment for posttraumatic stress disorder: Results of a preliminary randomized clinical trial. *Journal of Traumatic Stress, 24*(5), 591-595.
- Krakow, B., Hollifield, M., Johnston, L., Koss, M., Schrader, R., Warner, T. D., Tandberg, D., . . . Prince, H. (2001). Imagery rehearsal therapy for chronic nightmares in sexual assault survivors with posttraumatic stress disorder: a randomized controlled trial. *Journal of the American Medical Association, 286*(5), 537-45.
- Kubany, E.S., Hill, E.E., & Owens, J.A. (2003). Cognitive trauma therapy for battered women with PTSD: preliminary findings. *Journal of Traumatic Stress, 16*(1), 81-91.
- Kubany, E.S., Hill, E.E., Owens, J.A., Iannce-Spencer, C., McCaig, M.A., Tremayne, K.J., & Williams, P.L. (2004). Cognitive trauma therapy for battered women with PTSD (CTT-BW). *Journal of Consulting and Clinical Psychology, 72*(1), 3-18.
- Lindauer, R.J., Gersons, B.P., van Meijl, E.P., Blom, K., Carlier, I.V., Vrijlandt, I., & Olff, M. (2005). Effects of brief eclectic psychotherapy in patients with posttraumatic stress disorder: randomized clinical trial. *Journal of Traumatic Stress, 18*(3), 205-12.
- Maercker, A., Zoellner, T., Menning, H., Rabe, S., Karl, A. (2006). Dresden PTSD treatment study: randomized controlled trial of motor vehicle accident survivors. *BMC Psychiatry, 6*(1), 29-36.
- Markowitz, J.C., Petkova, E., Neria, Y., Van Meter P.E., Zhao, Y., Hembree, E., Lovell, K., . . . Marshall, R.D. (2015). Is exposure necessary? A randomized clinical trial of interpersonal psychotherapy for PTSD. *American Journal of Psychiatry, 172*(5), 430-440.
- Marks, I., Lovell, K., Noshirvani, H., Livanou, M., & Thrasher, S. (1998). Treatment of posttraumatic stress disorder by exposure and/or cognitive restructuring: a controlled study. *Archives of General Psychiatry, 55*(4), 317-25.
- Maxwell, K., Callahan, J.L., Holtz, P., Janis, B.M., Gerber, M.M., & Connor, D.R. (2015). Comparative study of group treatments for posttraumatic stress disorder. *Psychotherapy*.
- McDonagh, A., Friedman, M., McHugo, G., Ford, J., Sengupta, A., Mueser, K., Demment, C.C., ... Descamps, M. (2005). Randomized trial of cognitive-behavioral therapy for chronic posttraumatic stress disorder in adult female survivors of childhood sexual abuse. *Journal of Consulting and Clinical Psychology, 73*(3), 515-524.
- Mills, K.L., Teesson, M., Back, S.E., Brady, K.T., Baker, A.L., Hopwood, S., Sannibale, C., . . . Ewer, P.L. (2012). Integrated exposure-based therapy for co-occurring posttraumatic stress disorder and substance dependence: a randomized controlled trial. *Jama, 308*(7), 690-9.
- Monson, C.M., Schnurr, P.P., Resick, P.A., Friedman, M.J., Young-Xu, Y., & Stevens, S.P. (2006). Cognitive processing therapy for veterans with military-related posttraumatic stress disorder. *Journal of Consulting and Clinical Psychology, 74*(5), 898-907.
- Monson, C.M., Fredman, S.J., Macdonald, A., Pukay-Martin, N.D., Resick, P.A., & Schnurr, P.P. (2012). Effect of cognitive-behavioral couple therapy for PTSD: a randomized controlled trial. *Jama, 308*(7), 700-9.
- Mueser, K.T., Rosenberg, S.D., Xie, H., Jankowski, M.K., Bolton, E.E., Lu, W., Hamblen, J.L., . . . Wolfe, R. (2008). A randomized controlled trial of cognitive-behavioral treatment for posttraumatic stress disorder in severe mental illness. *Journal of Consulting and Clinical Psychology, 76*(2), 259-71.
- Mueser, K.T., Gottlieb, J.D., Xie, H., Lu, W., Yanos, P.T., Rosenberg, S.D., Silverstein, S.M., . . . McHugo, G.J. (2015). Evaluation of cognitive restructuring for post-traumatic stress disorder in people with severe mental illness. *The British Journal of Psychiatry: the Journal of Mental Science, 206*(6), 501-8.
- Nacasch, N., Tzur, D., Fostick, L., Dinstein, Y., Polliack, M., Zohar, J., Foa, E.B., . . . Huppert, J.D. (2011). Prolonged exposure therapy for combat- and terror-related posttraumatic stress disorder: A randomized control comparison with treatment as usual. *Journal of Clinical Psychiatry, 72*(9), 1174-1180.
- Neuner, F., Schauer, M., Klaschik, C., Karunakara, U., & Elbert, T. (2004). A comparison of narrative exposure therapy, supportive counseling, and psychoeducation for treating posttraumatic stress disorder in an African refugee settlement. *Journal of Consulting and Clinical Psychology, 72*(4), 579-587.
- Neuner, F., Ertl, V., Odenwald, M., Schauer, E., Elbert, T., & Onyut, P.L. (2008). Treatment of posttraumatic stress disorder by trained lay counselors in an African refugee settlement: A randomized controlled trial. *Journal of Consulting and Clinical Psychology, 76*(4), 686-694.
- Neuner, F., Kurreck, S., Ruf, M., Odenwald, M., Elbert, T., & Schauer, M. (2010). Can asylum-seekers with posttraumatic stress disorder be successfully treated? A randomized controlled pilot study. *Cognitive Behaviour Therapy, 39*(2), 81-91.
- Power, K., McGoldrick, T., Brown, K., Buchanan, R., Sharp, D., Swanson, V., & Karatzias, A. (2002). A controlled comparison of eye movement desensitization and reprocessing versus exposure plus cognitive restructuring versus waiting list in the treatment of post-traumatic stress disorder. *Clinical Psychology & Psychotherapy, 9*(5), 299-318.
- Resick, P.A., & Schnicke, M.K. (1992). Cognitive processing therapy for sexual assault victims. *Journal of Consulting and Clinical Psychology, 60*(5), 748-756.
- Resick, P.A., Nishith, P., Weaver, T.L., Astin, M.C., & Feuer, C.A. (2002). A comparison of cognitive-processing therapy with prolonged exposure and a waiting condition for the treatment of chronic posttraumatic stress disorder in female rape victims. *Journal of Consulting and Clinical Psychology, 70*(4), 867-879.
- Schnurr, P.P., Friedman, M.J., Foy, D.W., Shea, M.T., Hsieh, F.Y., Lavori, P.W., Glynn, S.M., . . . Bernardy, N. C. (2003). Randomized trial of trauma-focused group therapy for posttraumatic stress disorder: results from a department of veterans affairs cooperative study. *Archives of General Psychiatry, 60*(5), 481-9.
- Schnurr, P.P., Friedman, M.J., Engel, C.C., Foa, E.B., Shea, M.T., Chow, B.K., Resick, P.A., . . . Bernardy, N. (2007). Cognitive behavioral therapy for posttraumatic stress disorder in women: A randomized controlled trial. *Jama: the Journal of the American Medical Association, 297*(8), 820-830.
- Taylor, S., Thordarson, D.S., Maxfield, L., Fedoroff, I.C., Lovell, K., & Ogradniczuk, J. (2003). Comparative efficacy, speed, and adverse effects of three PTSD treatments: exposure therapy, EMDR, and relaxation training. *Journal of Consulting and Clinical Psychology, 71*(2), 330-338.
- van Emmerik, A.A.P., Kamphuis, J.H., & Emmelkamp, P.M.G. (2008). Treating acute stress disorder and posttraumatic stress disorder with cognitive behavioral therapy or structured writing therapy: A randomized controlled trial. *Psychotherapy and Psychosomatics, 77*(2), 93-100.
- Zang, Y., Hunt, N., & Cox, T. (2013). A randomised controlled pilot study: the effectiveness of narrative exposure therapy with adult survivors of the Sichuan earthquake. *BMC Psychiatry, 13*(1), 1.
- Zang, Y., Hunt, N., & Cox, T. (2014). Adapting narrative exposure therapy for Chinese earthquake survivors: a pilot randomised controlled feasibility study. *BMC Psychiatry, 14*(1), 1.

For further information, contact:
(360) 664-9800, institute@wsipp.wa.gov

Printed on 05-25-2022



Washington State Institute for Public Policy

The Washington State Legislature created the Washington State Institute for Public Policy in 1983. A Board of Directors—representing the legislature, the governor, and public universities—governs WSIPP and guides the development of all activities. WSIPP's mission is to carry out practical research, at legislative direction, on issues of importance to Washington State.