

## Cost sharing: (n) Copays for prescription drugs, adults with a chronic illness

Health Care: Health Care System Efficiency  
Literature review updated November 2015.

As part of WSIPP's research approach to identifying evidence-based programs and policies, WSIPP determines "what works" (and what does not work) to improve outcomes using an approach called meta-analysis. For detail on our methods, see our [Technical Documentation](#). At this time, WSIPP has not yet calculated benefits and costs for this topic.

Program Description: Evaluations of health care policies and programs often measure two broad types of outcomes: (1) those that reflect the health status of people (e.g., disease incidence) and (2) those that reflect health care system costs and utilization. Cost and utilization measures may or may not be an indication of health status or well-being.

The effect reported below is for increases in prescription drug copays (ranging from \$8 to \$23) in employer-sponsored health plans. The estimate is for patients taking medications for hypertension and high cholesterol (ACE inhibitors and statins).

### Meta-Analysis of Program Effects

Outcomes measured	No. of effect sizes	Treatment N	Adjusted effect size and standard error			Unadjusted effect size (random effects model)	
			ES	SE	Age	ES	p-value
Medication adherence	2	652	-0.602	0.118	30	-0.602	0.001

Meta-analysis is a statistical method to combine the results from separate studies on a program, policy, or topic in order to estimate its effect on an outcome. WSIPP systematically evaluates all credible evaluations we can locate on each topic. The outcomes measured are the types of program impacts that were measured in the research literature (for example, crime or educational attainment). Treatment N represents the total number of individuals or units in the treatment group across the included studies.

An effect size (ES) is a standard metric that summarizes the degree to which a program or policy affects a measured outcome. If the effect size is positive, the outcome increases. If the effect size is negative, the outcome decreases.

Adjusted effect sizes are used to calculate the benefits from our benefit cost model. WSIPP may adjust effect sizes based on methodological characteristics of the study. For example, we may adjust effect sizes when a study has a weak research design or when the program developer is involved in the research. The magnitude of these adjustments varies depending on the topic area.

WSIPP may also adjust the second ES measurement. Research shows the magnitude of some effect sizes decrease over time. For those effect sizes, we estimate outcome-based adjustments which we apply between the first time ES is estimated and the second time ES is estimated. We also report the unadjusted effect size to show the effect sizes before any adjustments have been made. More details about these adjustments can be found in our [Technical Documentation](#).

### Citations Used in the Meta-Analysis

Huskamp, H.A., Deverka, P.A., Epstein, A.M., Epstein, R.S., McGuigan, K.A., & Frank, R.G. (2003). The effect of incentive-based formularies on prescription-drug utilization and spending. *The New England Journal of Medicine*, 349(23), 2224-32.

For further information, contact:  
(360) 664-9800, [Institute@wsipp.wa.gov](mailto:Institute@wsipp.wa.gov)

Printed on 04-02-2024



## Washington State Institute for Public Policy

The Washington State Legislature created the Washington State Institute for Public Policy in 1983. A Board of Directors—representing the legislature, the governor, and public universities—governs WSIPP and guides the development of all activities. WSIPP's mission is to carry out practical research, at legislative direction, on issues of importance to Washington State.