

Preschool programs to create a healthy food environment and increase physical activity

Public Health & Prevention: School-based

Benefit-cost estimates updated December 2019. Literature review updated November 2015.

Current estimates replace old estimates. Numbers will change over time as a result of model inputs and monetization methods.

The WSIPP benefit-cost analysis examines, on an apples-to-apples basis, the monetary value of programs or policies to determine whether the benefits from the program exceed its costs. WSIPP's research approach to identifying evidence-based programs and policies has three main steps. First, we determine "what works" (and what does not work) to improve outcomes using a statistical technique called meta-analysis. Second, we calculate whether the benefits of a program exceed its costs. Third, we estimate the risk of investing in a program by testing the sensitivity of our results. For more detail on our methods, see our [Technical Documentation](#).

Program Description: A wide variety of interventions were used in the preschool programs that we reviewed. These included nutrition education for children and parents, additional time engaged in physical activity, and healthier snacks and drinks during the school day. The interventions were conducted in preschools serving children three to six years old.

Benefit-Cost Summary Statistics Per Participant

Benefits to:

Taxpayers	\$1	Benefit to cost ratio	(\$0.49)
Participants	\$0	Benefits minus costs	(\$393)
Others	\$1	Chance the program will produce	
Indirect	(\$132)	benefits greater than the costs	45 %
Total benefits	(\$129)		
Net program cost	(\$264)		
Benefits minus cost	(\$393)		

The estimates shown are present value, life cycle benefits and costs. All dollars are expressed in the base year chosen for this analysis (2018). The chance the benefits exceed the costs are derived from a Monte Carlo risk analysis. The details on this, as well as the economic discount rates and other relevant parameters are described in our [Technical Documentation](#).

Meta-Analysis of Program Effects

Outcomes measured	Treatment age	No. of effect sizes	Treatment N	Adjusted effect sizes and standard errors used in the benefit-cost analysis						Unadjusted effect size (random effects model)	
				First time ES is estimated			Second time ES is estimated			ES	p-value
				ES	SE	Age	ES	SE	Age		
Obesity	5	12	4490	-0.140	0.039	6	0.000	0.101	8	-0.140	0.001

Meta-analysis is a statistical method to combine the results from separate studies on a program, policy, or topic in order to estimate its effect on an outcome. WSIPP systematically evaluates all credible evaluations we can locate on each topic. The outcomes measured are the types of program impacts that were measured in the research literature (for example, crime or educational attainment). Treatment N represents the total number of individuals or units in the treatment group across the included studies.

An effect size (ES) is a standard metric that summarizes the degree to which a program or policy affects a measured outcome. If the effect size is positive, the outcome increases. If the effect size is negative, the outcome decreases.

Adjusted effect sizes are used to calculate the benefits from our benefit cost model. WSIPP may adjust effect sizes based on methodological characteristics of the study. For example, we may adjust effect sizes when a study has a weak research design or when the program developer is involved in the research. The magnitude of these adjustments varies depending on the topic area.

WSIPP may also adjust the second ES measurement. Research shows the magnitude of some effect sizes decrease over time. For those effect sizes, we estimate outcome-based adjustments which we apply between the first time ES is estimated and the second time ES is estimated. We also report the unadjusted effect size to show the effect sizes before any adjustments have been made. More details about these adjustments can be found in our [Technical Documentation](#).

Detailed Monetary Benefit Estimates Per Participant						
Affected outcome:	Resulting benefits: ¹	Benefits accrue to:				
		Taxpayers	Participants	Others ²	Indirect ³	Total
Obesity	Labor market earnings associated with obesity	\$0	\$0	\$0	\$0	\$0
Obesity	Health care associated with obesity	\$1	\$0	\$1	\$0	\$2
Obesity	Mortality associated with obesity	\$0	\$0	\$0	\$0	\$0
Program cost	Adjustment for deadweight cost of program	\$0	\$0	\$0	(\$132)	(\$132)
Totals		\$1	\$0	\$1	(\$132)	(\$129)

¹In addition to the outcomes measured in the meta-analysis table, WSIPP measures benefits and costs estimated from other outcomes associated with those reported in the evaluation literature. For example, empirical research demonstrates that high school graduation leads to reduced crime. These associated measures provide a more complete picture of the detailed costs and benefits of the program.

²"Others" includes benefits to people other than taxpayers and participants. Depending on the program, it could include reductions in crime victimization, the economic benefits from a more educated workforce, and the benefits from employer-paid health insurance.


³"Indirect benefits" includes estimates of the net changes in the value of a statistical life and net changes in the deadweight costs of taxation.

Detailed Annual Cost Estimates Per Participant				
	Annual cost	Year dollars	Summary	
Program costs	\$247	2014	Present value of net program costs (in 2018 dollars) (\$264)	
Comparison costs	\$0	2014	Cost range (+ or -) 20 %	

The programs in this analysis added 123 additional hours of physical activity or nutrition curriculum; we assume that these occur within a single school year. We assumed there was a maximum of 20 students per class and an adult-to-child ratio of 1:10. The per-student cost of the intervention was calculated by multiplying the per-student staff hours required by two staff members whose compensation were the average salary and benefits of a preschool teacher in 2014 in Washington State reported by the Bureau of Labor Statistics.


The figures shown are estimates of the costs to implement programs in Washington. The comparison group costs reflect either no treatment or treatment as usual, depending on how effect sizes were calculated in the meta-analysis. The cost range reported above reflects potential variation or uncertainty in the cost estimate; more detail can be found in our [Technical Documentation](#).

Benefits Minus Costs Over Time (Cumulative Discounted Dollars)




The graph above illustrates the estimated cumulative net benefits per-participant for the first fifty years beyond the initial investment in the program. We present these cash flows in discounted dollars. If the dollars are negative (bars below \$0 line), the cumulative benefits do not outweigh the cost of the program up to that point in time. The program breaks even when the dollars reach \$0. At this point, the total benefits to participants, taxpayers, and others, are equal to the cost of the program. If the dollars are above \$0, the benefits of the program exceed the initial investment.

Benefits by Perspective Over Time (Cumulative Discounted Dollars)



The graph above illustrates the breakdown of the estimated cumulative benefits (not including program costs) per-participant for the first fifty years beyond the initial investment in the program. These cash flows provide a breakdown of the classification of dollars over time into four perspectives: taxpayer, participant, others, and indirect. "Taxpayers" includes expected savings to government and expected increases in tax revenue. "Participants" includes expected increases in earnings and expenditures for items such as health care and college tuition. "Others" includes benefits to people other than taxpayers and participants. Depending on the program, it could include reductions in crime victimization, the economic benefits from a more educated workforce, and the benefits from employer-paid health insurance. "Indirect benefits" includes estimates of the changes in the value of a statistical life and changes in the deadweight costs of taxation. If a section of the bar is below the \$0 line, the program is creating a negative benefit, meaning a loss of value from that perspective.



The graph above focuses on the subset of estimated cumulative benefits that accrue to taxpayers. The cash flows are divided into the source of the value.

Citations Used in the Meta-Analysis

- Bayer, O., von Kries, R., Strauss, A., Mitschek, C., Toschke, A. M., Hose, A. et al. (2009). Short- and mid-term effects of a setting based prevention program to reduce obesity risk factors in children: a cluster-randomized trial. *Clinical Nutrition*, 28, 122-128.
- De Bock, F., Breitenstein, L., & Fischer, J. E. (2012). Positive impact of a pre-school-based nutritional intervention on children's fruit and vegetable intake: results of a cluster-randomized trial. *Public Health Nutrition*, 15(3), 466-475.
- Eliakim, A., Nemet, D., Balakirski, Y., & Epstein, Y. (2007). The effects of nutritional-physical activity school-based intervention on fatness and fitness in preschool children. *Journal of Pediatric Endocrinology & Metabolism*, 20(6), 711-718.
- Fitzgibbon, M.L., Stolley, M.R., Schiffer, L., Van Horn, L., KauferChristoffel, K., & Dyer, A. (2005). Two-year follow-up results for Hip-Hop to Health Jr.: a randomized controlled trial for overweight prevention in preschool minority children. *The Journal of pediatrics*, 146(5), 618-625.
- Fitzgibbon, M.L., Stolley, M.R., Schiffer, L., Van, H.L., Kauferchristoffel, K., Dyer, A. (2006). Hip-Hop to Health Jr. for Latino Preschool Children. *Obesity*, 14(9), 1616.
- Fitzgibbon, M.L., Stolley, M.R., Schiffer, L.A., Braunschweig, C.L., Gomez, S.L., Van, H.L., & Dyer, A.R. (2011). Hip-Hop to Health Jr. Obesity Prevention Effectiveness Trial: postintervention results. *Obesity*, 19(5), 994-1003.
- Jouret, B., Ahluwalia, N., Dupuy, M., Cristini, C., Negre-Pages, L., Grandjean, H., & Tauber, M. (2009). Prevention of overweight in preschool children: results of kindergarten-based interventions. *International Journal of Obesity*, 33(10), 1075-1083.
- Nemet, D., Geva, D., Pantanowitz, M., Igarria, N., Meckel, Y., & Eliakim, A. (2011). Health promotion intervention in Arab-Israeli kindergarten children. *Journal of Pediatric Endocrinology & Metabolism*, 24(11-12), 1001-1007.
- Nemet, D., Geva, D., & Eliakim, A. (2011). Health Promotion Intervention in Low Socioeconomic Kindergarten Children. *The Journal of Pediatrics*, 158(5), 796-801.
- Puder, JJ, Marques-Vidal, P, Schindler, C, Zahner, L, Niederer, I, Bürgli, F, Ebenegger, V, ... Kriemler, S. (2011). Effect of multidimensional lifestyle intervention on fitness and adiposity in predominantly migrant preschool children (Ballabeina): cluster randomised controlled trial. *BMJ*, 343, d6195.
- Reilly, J.J., Kelly, L., Montgomery, C., Williamson, A., Fisher, A., McColl, J.H., ... & Grant, S. (2006). Physical activity to prevent obesity in young children: cluster randomised controlled trial. *BMJ*, 333(7577), 1041-3.
- Zask, A., Adams, J.K., Brooks, L.O., & Hughes, D.F. (2012). Tooty Fruity Veggie: an obesity prevention intervention evaluation in Australian preschools. *Health Promotion Journal of Australia*, 23(1), 10-5.

For further information, contact:
(360) 664-9800, institute@wsipp.wa.gov

Printed on 07-05-2022



Washington State Institute for Public Policy

The Washington State Legislature created the Washington State Institute for Public Policy in 1983. A Board of Directors—representing the legislature, the governor, and public universities—governs WSIPP and guides the development of all activities. WSIPP's mission is to carry out practical research, at legislative direction, on issues of importance to Washington State.